

美定隆® 糖衣錠60毫克

(溴化吡啶斯狄明)

膽素脂酶抑制劑，抗重症肌無力

成份：

每錠含pyridostigmine bromide 60mg

賦型劑：

Maize Starch, Magnesium Stearate, Povidone, Pregelatinized Potato Starch, Aerosil, Talc, Acacia, Iron oxide red, Iron oxide yellow, Liquid Paraffin, Paraffin, Rice Starch, Sucrose

性能：

美定隆可抑制膽素脂酶而有效激活膽素的功能，作用溫和藥效長。美定隆特別適用於治療重症肌無力。其副作用輕微。

適應症：

重症肌無力。

劑量：

重症肌無力

每天2~4次，每次1~3錠

注意事項：

美定隆會加強 morphine 衍生物和 barbiturates 的作用。病人患有氣喘或痙攣性氣管炎時，使用美定隆要特別謹慎。

備註：

皮下注射或肌肉注射0.4mg atropine sulfate可中和擬副交感神經藥的作用。

包裝：

4-1000粒瓶裝

衛署藥製字第036008號

本藥須由醫師處方使用

請置於孩童取用不到之處

藥商：新加坡商美納里尼醫藥有限公司 台灣分公司

地址：台北市大安區敦化南路二段65號17樓

電話：02-2708-1863

製造廠：歐帕生技醫藥股份有限公司

廠址：新竹縣湖口鄉光復路1號



MENARINI

Mestinon® S.C. Tablets 60mg

(Pyridostigmine bromide)

Anticholinesterase, antimuscarinic

Composition

Pyridostigmine bromide: dimethylcarbamic ester of 1-methyl-3-hydroxypyridinium bromide.

Excipients

Maize Starch, Magnesium Stearate, Povidone, Pregelatinized Potato Starch, Aerosil, Talc, Acacia, Iron oxide red, Iron oxide yellow, Liquid Paraffin, Paraffin, Rice Starch, Sucrose

Properties

Mestinon has cholinergic properties; it is a cholinesterase inhibitor. It is characterized by the gentle onset, smooth course, comparatively long duration and gradual decline of its action.

Mestinon is particularly suitable for use as an antimuscarinic. The side effects of Mestinon are mild.

Indications/Dosage

Myasthenia gravis: 1-3 tablets twice to four times daily, or more in severe cases

Precautions

The action of morphine derivatives and barbiturates may be potentiated by Mestinon. Mestinon should be used with caution in patients with bronchial asthma or spastic bronchitis.

Note

Any parasympathomimetic effects can be counteracted by s.c. or i.m. injection of 0.4 mg atropine sulfate.

Packs

4-1000 Tablets/Bottle



MENARINI

Manufactured by
AUPA BIOPHARM CO., LTD. for
A. Menarini Singapore Pte. Ltd., Taiwan Branch